

ON THE DISTRIBUTION OF THE FAREY SEQUENCE WITH ODD DENOMINATORS

FLORIN P. BOCA, CRISTIAN COBELI AND ALEXANDRU ZAHARESCU

1. INTRODUCTION AND STATEMENT OF RESULTS

Given a positive integer Q , we denote by $\mathcal{F}_Q$ the set of irreducible rational fractions in $(0, 1]$ whose denominators do not exceed Q . That is,

$$\mathcal{F}_Q = \{a/q; 1 \leq a \leq q \leq Q, \gcd(a, q) = 1\}.$$

Problems concerning the distribution of Farey fractions have been studied in the 20's by Franel and Landau ([6],[15]) and more recently in [1], [2], [3], [4], [7], [8], [9], [10], [11], [13], [14].

It is well-known that

$$N_Q = \#\mathcal{F}_Q = 6Q^2/\pi^2 + O(Q \log Q).$$

We denote by $\mathcal{F}_Q^<$ the set of pairs (γ, γ') of consecutive elements in $\mathcal{F}_Q$.

In this paper we are concerned with the set

$$\mathcal{F}_{Q,\text{odd}} = \{a/q \in \mathcal{F}_Q; q \text{ odd}\}$$

of Farey fractions of order Q with odd denominators. For instance,

$$\begin{aligned} \mathcal{F}_8 &= \left\{ \frac{1}{8}, \frac{1}{7}, \frac{1}{6}, \frac{1}{5}, \frac{1}{4}, \frac{2}{7}, \frac{1}{3}, \frac{3}{8}, \frac{2}{5}, \frac{3}{7}, \frac{1}{2}, \frac{4}{7}, \frac{3}{5}, \frac{5}{8}, \frac{2}{3}, \frac{5}{7}, \frac{3}{4}, \frac{5}{6}, \frac{6}{7}, \frac{7}{8}, 1 \right\} \quad \text{and} \\ \mathcal{F}_{8,\text{odd}} &= \left\{ \frac{1}{7}, \frac{1}{5}, \frac{2}{7}, \frac{1}{3}, \frac{2}{5}, \frac{3}{7}, \frac{4}{5}, \frac{3}{3}, \frac{2}{7}, \frac{5}{5}, \frac{4}{3}, \frac{6}{7}, 1 \right\}. \end{aligned}$$

The set of pairs (γ, γ') of consecutive elements in $\mathcal{F}_{Q,\text{odd}}$ is denoted by $\mathcal{F}_{Q,\text{odd}}^<$. It is not hard to prove (see [11]) that

$$(1.1) \quad N_{Q,\text{odd}} = \#\mathcal{F}_{Q,\text{odd}} = 2Q^2/\pi^2 + O(Q \log Q).$$

It is well-known that $\Delta(\gamma, \gamma') := a'q - aq' = 1$ whenever $\gamma = \frac{a}{q} < \frac{a'}{q'} = \gamma'$ are consecutive elements in $\mathcal{F}_Q$. This certainly fails when $\gamma < \gamma'$ are consecutive in $\mathcal{F}_{Q,\text{odd}}$. A first step in the study of the distribution of the values of $\Delta(\gamma, \gamma')$ for pairs (γ, γ') of consecutive fractions in $\mathcal{F}_{Q,\text{odd}}$ was undertaken by A. Haynes in [11]. He proved that if one denotes

$$N_{Q,\text{odd}}(k) = \#\{\gamma < \gamma' \text{ successive in } \mathcal{F}_{Q,\text{odd}}; \Delta(\gamma, \gamma') = k\},$$

then the asymptotic frequency

$$\rho_{\text{odd}}(k) = \lim_{Q \rightarrow \infty} \frac{N_{Q,\text{odd}}(k)}{N_{Q,\text{odd}}}$$

Research partially supported by ANSTI grant C6189/2000.

exists, and is expressed as

$$\rho_{\text{odd}}(k) = \frac{4}{k(k+1)(k+2)}, \quad k \in \mathbb{N}^*.$$

This can be written as

$$(1.2) \quad \rho_{\text{odd}}(k) = \begin{cases} \text{Area}(\mathcal{T}_k) & \text{if } k \geq 2 \\ \frac{1}{2} + \text{Area}(\mathcal{T}_1) & \text{if } k = 1, \end{cases}$$

where (as in [3]) we denote $\mathcal{T}_k = \{(x, y) \in \mathcal{T} ; \lfloor \frac{1+x}{y} \rfloor = k\}$, $k \in \mathbb{N}^*$, and $\mathcal{T} = \{(x, y) \in [0, 1] ; x + y > 1\}$.

In this note we study, for fixed $h \geq 1$, the distribution of consecutive elements $\gamma_i < \gamma_{i+1} < \dots < \gamma_{i+h}$ in $\mathcal{F}_{Q, \text{odd}}$, and compute the probability that such an $(h+1)$ -tuple satisfies $\Delta(\gamma_i, \gamma_{i+1}) = \Delta_1, \dots, \Delta(\gamma_{i+h-1}, \gamma_{i+h}) = \Delta_h$. More precisely, we prove that if one denotes

$$N_{Q, \text{odd}}(\Delta_1, \dots, \Delta_h) = \# \left\{ i; \gamma_i < \gamma_{i+1} < \dots < \gamma_{i+h} \text{ consecutive in } \mathcal{F}_{Q, \text{odd}} \right. \\ \left. \Delta(\gamma_{i+j-1}, \gamma_{i+j}) = \Delta_j, \ j = 1, \dots, h \right\},$$

then

$$\rho_{\text{odd}}(\Delta_1, \dots, \Delta_h) = \lim_{Q \rightarrow \infty} \frac{N_{Q, \text{odd}}(\Delta_1, \dots, \Delta_h)}{N_{Q, \text{odd}}}$$

exists for all $h \geq 2$, and give an explicit formula for it.

To state the main result, we shall employ the area-preserving transformation T of $\mathcal{T}$, introduced in [3] and defined by

$$(1.3) \quad T(x, y) = \left(y, \left\lfloor \frac{1+x}{y} \right\rfloor y - x \right).$$

We denote

$$\mathcal{T}_{k_1, \dots, k_h} = \mathcal{T}_{k_1} \cap T^{-1} \mathcal{T}_{k_2} \cap \dots \cap T^{-h+1} \mathcal{T}_{k_h}.$$

We notice that if $\gamma = \frac{a}{q} < \gamma' = \frac{a'}{q'} < \gamma'' = \frac{a''}{q''}$ are consecutive elements in $\mathcal{F}_Q$, then $T(\frac{a}{q}, \frac{a'}{q'}) = (\frac{a'}{q'}, \frac{a''}{q''})$. Moreover, if we set $\kappa(x, y) = \lfloor \frac{1+x}{y} \rfloor$, then the positive integer $\kappa(\frac{a}{q}, \frac{a'}{q'}) = \lfloor \frac{Q+a}{q'} \rfloor$ coincides with the index $\nu_Q(\gamma)$ of the Farey fraction γ in $\mathcal{F}_Q$ considered in [10].

It will be worthwhile to consider the tree $\mathfrak{T}_h$ defined by the following properties:


- vertices are labeled by O and E ;
- the starting vertex $\star$ is labeled by O ;
- there is exactly one edge starting from an E vertex, and such an edge always ends into an O vertex;
- there are exactly two edges starting from an O vertex, and they end (respectively) into an E vertex and into an O vertex;
- the number of O vertices (besides $\star$) on any path that originates at $\star$ is equal to h .

See Figure 1.

We also consider the set $\mathfrak{L}_h$ of labeled paths

$$w = \left(\star = O \xrightarrow{k_1} v_1 \xrightarrow{k_2} v_2 \xrightarrow{k_3} \dots \xrightarrow{k_{|w|}} v_{|w|} \right), \quad k_j \in \mathbb{N}^*.$$

on the tree $\mathfrak{T}_h$ that start at $\star$ and pass through $h+1$ vertices labeled by O (including $\star$). That is, $\#\{j ; v_j = O\} = h$. We set $o(O) = \text{odd}$, $o(E) = \text{even}$.


$$\begin{aligned} c_{OE}(\star = O \xrightarrow{k_1} E \xrightarrow{k_2} O) &= k_1, & c_{OE}(\star = O \xrightarrow{k_1} O) &= \emptyset, \\ c_{\Delta_1}(\star = O \xrightarrow{k_1} E \xrightarrow{k_2} O) &= \Delta_1, & c_{\Delta_1}(\star = O \xrightarrow{k_1} O) &= \emptyset, \end{aligned}$$
$$c_{OE}(w) = \begin{cases} (c_{OE}(w'), k) & \text{if } w'' = O \xrightarrow{k} E \xrightarrow{l} O \\ c_{OE}(w') & \text{if } w'' = O \xrightarrow{k} O, \end{cases}$$

$$c_{(\Delta_1, \dots, \Delta_{h+1})}(w) = \begin{cases} (c_{\Delta}(w'), \Delta_{h+1}) & \text{if } w'' = O \xrightarrow{k} E \xrightarrow{l} O \\ c_{\Delta}(w') & \text{if } w'' = O \xrightarrow{k} O. \end{cases}$$
$$\star = O \xrightarrow{k_1} E \xrightarrow{k_2} O \xrightarrow{k_3} O \xrightarrow{k_4} E \xrightarrow{k_5} O \xrightarrow{k_6} O \xrightarrow{k_7} E \xrightarrow{k_8} O$$
$$c_{OE}(w) = (k_1, k_4, k_7) \quad \text{and} \quad c_{(\Delta_1, \dots, \Delta_5)}(w) = (\Delta_1, \Delta_3, \Delta_5).$$
$$v_0 = \star = O \xrightarrow{k_1} v_1 \xrightarrow{k_2} \cdots \xrightarrow{k_{|w|}} v_{|w|}$$

such that $c_{OE}(w) = c_\Delta(w)$, and such that k_j is even whenever it occurs as $E \xrightarrow{k_j} O \text{ --- } E$ or as $O \xrightarrow{k_j} O \text{ --- } O$, and respectively odd whenever it occurs as $E \xrightarrow{k_j} O \text{ --- } O$ or as $O \xrightarrow{k_j} O \text{ --- } E$.

Having established this notation, we may state our main result.

Theorem 1.1. *Let $h \geq 1$, and let $\Delta = (\Delta_1, \dots, \Delta_h) \in (\mathbb{N}^*)^h$. Then*

$$\rho_{Q,\text{odd}}(\Delta) := \frac{N_{Q,\text{odd}}(\Delta_1, \dots, \Delta_h)}{N_{Q,\text{odd}}} = \rho_{\text{odd}}(\Delta) + O_h\left(\frac{\log^2 Q}{Q}\right)$$

as $Q \rightarrow \infty$, where

$$(1.4) \quad \rho_{\text{odd}}(\Delta) = \sum_{w \in \mathfrak{L}_h \cap \mathfrak{S}_\Delta} \text{Area}(\mathcal{T}_{k_1, \dots, k_{|w|-1}}).$$

For $h = 1$, this gives

$$\rho_{\text{odd}}(\Delta_1) = \begin{cases} \sum_{k_1} \text{Area}(\mathcal{T}_{k_1}) + \text{Area}(\mathcal{T}_1) = \frac{1}{2} + \text{Area}(\mathcal{T}_1) & \text{if } \Delta_1 = 1 \\ \sum_{k_2} \text{Area}(\mathcal{T}_{\Delta_1} \cap T^{-1}\mathcal{T}_{k_2}) = \text{Area}(\mathcal{T}_{\Delta_1}) & \text{if } \Delta_1 \geq 2, \end{cases}$$

that is, the aforementioned result of Haynes.

For $h = 2$, we obtain the following

Corollary 1.2. *$\rho_{Q,\text{odd}}(\Delta_1, \Delta_2)$ tends to $\rho_{\text{odd}}(\Delta_1, \Delta_2)$ for any $\Delta_1, \Delta_2 \in \mathbb{N}^*$ as $Q \rightarrow \infty$. Moreover, we have:*

$$(i) \quad \rho_{\text{odd}}(1, 1) = \sum_{k_1 \text{ even}} \text{Area}(\mathcal{T}_{k_1}) + \sum_{k_1 \text{ odd}} \text{Area}(\mathcal{T}_{k_1, 1}) + \sum_{k_2 \text{ odd}} \text{Area}(\mathcal{T}_{1, k_2}) \\ + \sum_{k_2 \text{ even}} \text{Area}(\mathcal{T}_{1, k_2, 1}).$$

(ii) *if $\Delta_2 \geq 2$, then*

$$\rho_{\text{odd}}(1, \Delta_2) = \sum_{k_1 \text{ odd}} \text{Area}(\mathcal{T}_{k_1, \Delta_2}) + \sum_{k_2 \text{ even}} \text{Area}(\mathcal{T}_{1, k_2, \Delta_2}).$$

(iii) *if $\Delta_1 \geq 2$, then*

$$\rho_{\text{odd}}(\Delta_1, 1) = \sum_{k_2 \text{ odd}} \text{Area}(\mathcal{T}_{\Delta_1, k_2}) + \sum_{k_2 \text{ even}} \text{Area}(\mathcal{T}_{\Delta_1, k_2, 1}).$$

(iv) *if $\min(\Delta_1, \Delta_2) \geq 2$, then*

$$\rho_{\text{odd}}(\Delta_1, \Delta_2) = \sum_{k_2 \text{ even}} \text{Area}(\mathcal{T}_{\Delta_1, k_2, \Delta_2}).$$

Actually, it follows from Lemma 3.4 and Remark 3.5 that all sums in (ii), (iii) and (iv) are finite.

In this kind of situation, one can give a short interval version of Theorem 1.1. For each interval $I \subseteq [0, 1]$ and for each $\Delta = (\Delta_1, \dots, \Delta_h) \in (\mathbb{N}^*)^h$, let

$$\begin{aligned} N_{Q,\text{odd}}^I &= \#\{\gamma_0 < \dots < \gamma_h \text{ consecutive in } \mathcal{F}_{Q,\text{odd}}; \gamma_0 \in I\} \\ &= 2|I|Q^2/\pi^2 + O(Q \log Q), \\ N_{Q,\text{odd}}^I(\Delta) &= \#\left\{i; \gamma_i \in I, \gamma_i < \gamma_{i+1} < \dots < \gamma_{i+h} \text{ consecutive in } \mathcal{F}_{Q,\text{odd}} \right. \\ &\quad \left. \Delta(\gamma_{i+j-1}, \gamma_{i+j}) = \Delta_j, j = 1, \dots, h \right\}. \end{aligned}$$

Then the following result holds.

Theorem 1.3. *Let $h \geq 1$, and assume that $\Delta = (\Delta_1, \dots, \Delta_h) \in (\mathbb{N}^*)^h$ is such that only finitely many non-vanishing terms appear on the right-hand side of (1.4). Then for any interval $I \subseteq [0, 1]$ we have*

$$\rho_{Q,\text{odd}}^I(\Delta) := \frac{N_{Q,\text{odd}}^I(\Delta)}{N_{Q,\text{odd}}^I} = \rho_{\text{odd}}(\Delta) + O_{h,\varepsilon}(Q^{-1/2+\varepsilon})$$

for every $\varepsilon > 0$.

The main techniques of a proof involve the basic properties of Farey fractions, the transformation T from (1.3), and estimates of Weil type for Kloosterman sums (see [16], [12], [5]).

2. REDUCTION OF $N_{Q,\text{odd}}(\Delta_1, \dots, \Delta_h)$

We set throughout

$$\mathbb{Z}_{\text{pr}}^2 = \{(a, b) \in \mathbb{Z}^2; \gcd(a, b) = 1\},$$

and denote for any subset Ω of $\mathbb{R}^2$ and $Q \in \mathbb{N}^*$

$$\begin{aligned} \partial\Omega &= \text{the boundary of } \Omega, & Q\Omega &= \{(Qx, Qy); (x, y) \in \Omega\}, \\ M(\Omega) &= \#(\Omega \cap \mathbb{Z}^2), & M_{\text{odd}}(\Omega) &= \#\{(x, y) \in \Omega \cap \mathbb{Z}^2; x \text{ odd}\}, \\ M_{\text{even}}(\Omega) &= \#\{(x, y) \in \Omega \cap \mathbb{Z}^2; x \text{ even}\} = M(\Omega) - M_{\text{odd}}(\Omega), \\ N(\Omega) &= \#(\Omega \cap \mathbb{Z}_{\text{pr}}^2), & N_{\text{odd}}(\Omega) &= \#\{(x, y) \in \Omega \cap \mathbb{Z}_{\text{pr}}^2; x \text{ odd}\}, \\ N_{\text{even}}(\Omega) &= N(\Omega) - N_{\text{odd}}(\Omega) = \#\{(x, y) \in \Omega \cap \mathbb{Z}_{\text{pr}}^2; x \text{ even}\}, \\ N_{\text{odd,odd}}(\Omega) &= \#\{(x, y) \in \Omega \cap \mathbb{Z}_{\text{pr}}^2; x \text{ odd}, y \text{ odd}\}, \\ N_{\text{odd,even}}(\Omega) &= \#\{(x, y) \in \Omega \cap \mathbb{Z}_{\text{pr}}^2; x \text{ odd}, y \text{ even}\}, \\ N_{\text{even,odd}}(\Omega) &= \#\{(x, y) \in \Omega \cap \mathbb{Z}_{\text{pr}}^2; x \text{ even}, y \text{ odd}\}. \end{aligned}$$

If $\gamma_{i_0} = \frac{a_{i_0}}{q_{i_0}} < \gamma_{i_0+1} = \frac{a_{i_0+1}}{q_{i_0+1}} < \dots < \gamma_{i_0+h} = \frac{a_{i_0+h}}{q_{i_0+h}}$ are consecutive in $\mathcal{F}_Q$, then (cf. [3])

$$\left(\frac{q_{i_0+r}}{Q}, \frac{q_{i_0+r+1}}{Q}\right) = T^r\left(\frac{q_{i_0}}{Q}, \frac{q_{i_0+1}}{Q}\right).$$

There is a one-to-one correspondence between $\mathbb{Z}_{\text{pr}}^2 \cap Q\mathcal{T}_{k_1, \dots, k_r}$ and the set $\mathcal{F}_{Q, k_1, \dots, k_r}$ of consecutive elements $\gamma_0 < \gamma_1 < \dots < \gamma_r$ in $\mathcal{F}_Q$ with $\nu_Q(\gamma_{j-1}) = k_j$, $j = 1, \dots, r$, given by

$$(q_0, q_1) \mapsto (\gamma_0, \gamma_1, \dots, \gamma_r),$$

where (γ_0, γ_1) is the unique pair in $\mathcal{F}_Q^<$ with denominators q_0 and q_1 , and (γ_j, γ_{j+1}) the unique pair in $\mathcal{F}_Q^<$ with denominators $QT^j(\frac{q_0}{Q}, \frac{q_1}{Q})$, $j = 1, \dots, r$. This also shows that the set

$$\mathcal{F}_{Q, k_1, \dots, k_r}^{\text{odd, odd/even}} = \{(\gamma_0, \dots, \gamma_r) \in \mathcal{F}_{Q, k_1, \dots, k_r}; q_0 \text{ odd, } q_1 \text{ odd/even}\}$$

has cardinality $N_{\text{odd, odd/even}}(QT_{k_1, \dots, k_r})$.

Suppose that $\gamma = \frac{a}{q} < \frac{a''}{q''} = \gamma''$ are two consecutive elements in $\mathcal{F}_{Q, \text{odd}}$, and that $\Delta(\gamma, \gamma'') = a''q - aq'' > 1$. Since two fractions with even denominators cannot occur as consecutive elements in $\mathcal{F}_Q$, it follows that there is precisely one fraction $\gamma' = \frac{a'}{q'}$ in $\mathcal{F}_Q$ such that $\gamma < \gamma' < \gamma''$ are consecutive in $\mathcal{F}_Q$. One readily finds that (see for example [11, p. 4])

$$(2.1) \quad \Delta(\gamma, \gamma'') = \nu_Q(\gamma) = \left\lceil \frac{Q+q}{q'} \right\rceil = \kappa\left(\frac{q}{Q}, \frac{q'}{Q}\right).$$

To summarize, suppose that $\gamma < \gamma' < \gamma'' < \gamma''' < \gamma^{IV}$ are consecutive in $\mathcal{F}_Q$, and that q is odd. Denote by $q, q', \dots, q^{IV}$, respectively, the denominator of $\gamma, \gamma', \dots, \gamma^{IV}$. Denote also

$$k_j = k_j(q, q') = \kappa\left(T^{j-1}\left(\frac{q}{Q}, \frac{q'}{Q}\right)\right), \quad j \geq 1.$$

Then $q'' = k_1 q' - q$, $q''' = k_2 q'' - q'$ etc. The following situations may occur.

- (O) q' is odd and thus $\Delta(\gamma, \gamma') = 1$. Next, it could be *either* that
 - (OO) q'' is odd (if k_1 is even), in which case $(\gamma', \gamma'') \in \mathcal{F}_{Q, \text{odd}}^<$ and $\Delta(\gamma', \gamma'') = 1$, or that
 - (OEO) q'' is even (if k_1 is odd), in which case $q''' = k_2 q'' - q'$ is odd, $(\gamma', \gamma''') \in \mathcal{F}_{Q, \text{odd}}^<$, and $\Delta(\gamma', \gamma''') = k_2$.
- (E) q' is even, thus q'' is odd, $(\gamma, \gamma'') \in \mathcal{F}_{Q, \text{odd}}^<$ and $\Delta(\gamma, \gamma'') = k_1$. Next, we have *either* that
 - (EOO) q''' is odd (if k_2 is odd), in which case $(\gamma'', \gamma''') \in \mathcal{F}_{Q, \text{odd}}^<$ and $\Delta(\gamma'', \gamma''') = 1$, or that
 - (EOEO) q''' is even (if k_2 is even), in which case $q^{IV} = k_3 q''' - q''$ will also be odd, $(\gamma'', \gamma^{IV}) \in \mathcal{F}_{Q, \text{odd}}^<$ and $\Delta(\gamma'', \gamma^{IV}) = k_3$.

This suggests that one may express $N_{Q, \text{odd}}(\Delta_1, \dots, \Delta_h)$ for any $h \geq 1$ by an inductive procedure. Note first that

$$N_{Q, \text{odd}}(\Delta_1) = \begin{cases} \sum_{k_1} N_{\text{odd, odd}}(QT_{k_1}) + \sum_{k_2} N_{\text{odd, even}}(QT_{1, k_2}) \\ \quad = N_{\text{odd, odd}}(QT) + N_{\text{odd, even}}(QT_{\Delta_1}) & \text{if } \Delta_1 = 1 \\ \sum_{k_2} N_{\text{odd, even}}(QT_{\Delta_1, k_2}) = N_{\text{odd, even}}(QT_{\Delta_1}) & \text{if } \Delta_1 \geq 2. \end{cases}$$

One may also express $N_{Q,\text{odd}}(\Delta_1, \Delta_2)$ as

$$\left\{ \begin{array}{ll} \sum_{k_1 \text{ even}} N_{\text{odd,odd}}(Q\mathcal{T}_{k_1}) + \sum_{k_1 \text{ odd}} N_{\text{odd,odd}}(Q\mathcal{T}_{k_1,1}) \\ \quad + \sum_{k_2 \text{ odd}} N_{\text{odd,even}}(Q\mathcal{T}_{1,k_2}) + \sum_{k_2 \text{ even}} N_{\text{odd,even}}(Q\mathcal{T}_{1,k_2,1}) & \text{if } \Delta_1 = \Delta_2 = 1 \\ \sum_{k_1 \text{ odd}} N_{\text{odd,odd}}(Q\mathcal{T}_{k_1,\Delta_2}) + \sum_{k_2 \text{ even}} N_{\text{odd,even}}(Q\mathcal{T}_{1,k_2,\Delta_2}) & \text{if } \Delta_1 = 1, \Delta_2 \geq 2 \\ \sum_{k_2 \text{ odd}} N_{\text{odd,even}}(Q\mathcal{T}_{\Delta_1,k_2}) + \sum_{k_2 \text{ even}} N_{\text{odd,even}}(Q\mathcal{T}_{\Delta_1,k_2,1}) & \text{if } \Delta_1 \geq 2, \Delta_2 = 1 \\ \sum_{k_2 \text{ even}} N_{\text{odd,even}}(Q\mathcal{T}_{\Delta_1,k_2,\Delta_2}) & \text{if } \Delta_1, \Delta_2 \geq 2. \end{array} \right.$$

For $h \geq 2$, $\rho_{Q,\text{odd}}(\Delta_1, \dots, \Delta_h)$ is expressed in

Proposition 2.1. *Assume that $h \geq 2$ and $\Delta = (\Delta_1, \dots, \Delta_h) \in (\mathbb{N}^*)^h$. Then*

$$(2.2) \quad \rho_{Q,\text{odd}}(\Delta) = \frac{1}{N_{Q,\text{odd}}} \sum_{w \in \mathfrak{L}_h \cap \mathfrak{S}_\Delta} N_{\text{odd},o(v_1)}(Q\mathcal{T}_{k_1, \dots, k_{|w|-1}}).$$

3. ESTIMATING $N_{\text{odd,odd}}(\Omega)$ AND $N_{\text{odd,even}}(\Omega)$

For a bounded region Ω in $\mathbb{R}^2$ with rectifiable boundary and a function f defined on Ω , we set

$$\begin{aligned} S_f(\Omega) &= \sum_{(a,b) \in \Omega \cap \mathbb{Z}^2} f(a,b), & S'_f(\Omega) &= \sum_{(a,b) \in \Omega \cap \mathbb{Z}_{\text{pr}}^2} f(a,b), \\ S_{f,\text{odd/even}}(\Omega) &= \sum_{\substack{(a,b) \in \Omega \cap \mathbb{Z}^2 \\ a \text{ odd/even}}} f(a,b), & S'_{f,\text{odd/even}} &= \sum_{\substack{(a,b) \in \Omega \cap \mathbb{Z}_{\text{pr}}^2 \\ a \text{ odd/even}}} f(a,b), \\ S'_{f,\text{odd,odd/even}}(\Omega) &= \sum_{\substack{(a,b) \in \Omega \cap \mathbb{Z}_{\text{pr}}^2 \\ a \text{ odd}, b \text{ odd/even}}} f(a,b), \\ \|Df\|_{L^\infty(\Omega)} &= \sup_{(x,y) \in \Omega} \left(\left| \frac{\partial f}{\partial x}(x,y) \right| + \left| \frac{\partial f}{\partial y}(x,y) \right| \right). \end{aligned}$$

Lemma 3.1. *Let $R_1, R_2 > 0$, and let $R \geq \min(R_1, R_2)$. Then for any region $\Omega \subseteq [0, R_1] \times [0, R_2]$ and any function f which is C^1 on Ω , we have*

$$\begin{aligned} \text{(i)} \quad S'_{f,\text{odd}}(\Omega) &= \frac{4}{\pi^2} \iint_{\Omega} f(x,y) dx dy + O(A_{f,R,\Omega}). \\ \text{(ii)} \quad S'_{f,\text{odd,odd/even}}(\Omega) &= \frac{2}{\pi^2} \iint_{\Omega} f(x,y) dx dy + O(A_{f,R,\Omega}). \\ \text{(iii)} \quad S'_{f,\text{even,odd}}(\Omega) &= \frac{2}{\pi^2} \iint_{\Omega} f(x,y) dx dy + O(A_{f,R,\Omega}), \end{aligned}$$

where

$$A_{f,R,\Omega} = \frac{\|f\|_{L^1(\Omega)}}{R} + \|Df\|_{L^\infty(\Omega)} \text{Area}(\Omega) \log R \\ + \|f\|_{L^\infty(\Omega)} (R + \text{length}(\partial\Omega) \log R).$$

Proof. (i) It is well known (see e.g. [3, Lemma 1]) that

$$S_f(\Omega) = \iint_{\Omega} f(x, y) dx dy + O(B_{f,\Omega}),$$

where

$$B_{f,\Omega} = \|Df\|_{L^\infty(\Omega)} \text{Area}(\Omega) + \|f\|_{L^\infty(\Omega)} (1 + \text{length}(\partial\Omega)).$$

Denoting $\Omega' = \{(x/2, y); (x, y) \in \Omega\}$, we have that $S_{f,\text{even}}(\Omega)$ - and eventually $S_{f,\text{odd}}(\Omega)$ can be expressed as

$$(3.1) \quad \sum_{(a,b) \in \Omega' \cap \mathbb{Z}^2} f(2a, b) = \iint_{\Omega'} f(2x, y) dx dy + O(B_{f,\Omega'}) \\ = \frac{1}{2} \iint_{\Omega} f(x, y) dx dy + O(B_{f,\Omega}).$$

We now proceed to estimate $S'_{f,\text{odd}}(\Omega)$, which is written as

$$(3.2) \quad \sum_{\substack{(a,b) \in \Omega \\ a \text{ odd}}} f(a, b) - \sum_{\substack{(a,b) \in \Omega/3 \\ a \text{ odd}}} f(a, b) - \sum_{\substack{(a,b) \in \Omega/5 \\ a \text{ odd}}} f(a, b) - \dots \\ = \sum_{\substack{1 \leq n \leq R \\ n \text{ odd}}} \mu(n) \sum_{\substack{(a,b) \in \Omega/n \\ a \text{ odd}}} f(na, nb).$$

The inner sum in (3.2) is expressed by means of (3.1) as

$$(3.3) \quad \frac{1}{2} \iint_{\Omega/n} f(nx, ny) dx dy \\ + O\left(\frac{\|Df\|_{L^\infty(\Omega)} \text{Area}(\Omega)}{n} + \|f\|_{L^\infty(\Omega)} \left(1 + \frac{\text{length}(\partial\Omega)}{n}\right)\right).$$

Changing (nx, ny) to (x, y) in the double integral above and summing over n , we infer from (3.2) and (3.3) that

$$S'_{f,\text{odd}}(\Omega) = \frac{1}{2} \sum_{\substack{1 \leq n \leq R \\ n \text{ odd}}} \frac{\mu(n)}{n^2} \iint_{\Omega} f(x, y) dx dy + O(\|Df\|_{L^\infty(\Omega)} \text{Area}(\Omega) \log R) \\ + O\left(\|f\|_{L^\infty(\Omega)} (R + \text{length}(\partial\Omega) \log R)\right).$$

The equality (i) now follows from

$$\sum_{\substack{1 \leq n \leq R \\ n \text{ odd}}} \frac{\mu(n)}{n^2} = \frac{8}{\pi^2} + O\left(\frac{1}{R}\right).$$

The equality (ii) follows by combining (i) with

$$S'_{f,\text{odd},\text{even}}(\Omega) = \sum_{\substack{(a,b) \in \Omega'' \cap \mathbb{Z}_{\text{pr}}^2 \\ a \text{ odd}}} f(a, 2b),$$

where we set $\Omega'' = \{(x, y/2); (x, y) \in \Omega\}$, and then using

$$\iint_{\Omega''} f(x, 2y) dx dy = \frac{1}{2} \iint_{\Omega} f(x, y) dx dy.$$

The equality (iii) now follows from symmetry. $\square$

We need the following improvement of Lemma 1 in [11].

Corollary 3.2. *Let $R_1, R_2 > 0$, and let $R \geq \min(R_1, R_2)$. Then for any region $\Omega \subseteq [0, R_1] \times [0, R_2]$ with rectifiable boundary, we have*

- (i) $N_{\text{odd}}(\Omega) = 4\text{Area}(\Omega)/\pi^2 + O(C_{R,\Omega})$,
- (ii) $N_{\text{even}}(\Omega) = 2\text{Area}(\Omega)/\pi^2 + O(C_{R,\Omega})$,
- (iii) $N_{\text{odd},\text{even}}(\Omega) = 2\text{Area}(\Omega)/\pi^2 + O(C_{R,\Omega})$,
- (iv) $N_{\text{odd},\text{odd}}(\Omega) = 2\text{Area}(\Omega)/\pi^2 + O(C_{R,\Omega})$,
- (v) $N_{\text{even},\text{odd}}(\Omega) = 2\text{Area}(\Omega)/\pi^2 + O(C_{R,\Omega})$,

where

$$C_{R,\Omega} = \text{Area}(\Omega)/R + R + \text{length}(\partial\Omega) \log R.$$

The following lemma is contained in [3]. We enclose the proof for the reader's convenience.

Lemma 3.3. *For any integers $k_1, \dots, k_r \geq 1$, the set $\mathcal{T}_{k_1, \dots, k_r}$ is a convex polygon.*

Proof. If for $(x, y) \in \mathbb{R}^2$ we define $L_0(x, y) = x$, $L_1(x, y) = y$, and $L_{i+1}(x, y) = k_i L_i(x, y) - L_{i-1}(x, y)$ for $i \geq 1$, then $\mathcal{T}_{k_1, \dots, k_r}$ is defined by the following inequalities:

$$\begin{cases} 1 \geq L_0(x, y), L_1(x, y), \dots, L_{r+1}(x, y) > 0, \\ L_0(x, y) + L_1(x, y), L_1(x, y) + L_2(x, y), \dots, L_r(x, y) + L_{r+1}(x, y) > 1. \end{cases}$$

Because $L_0, L_1, \dots, L_{r+1}$ are linear functions, the set $\mathcal{T}_{k_1, \dots, k_r}$ is the intersection of finitely many convex polygons. $\square$

Lemma 3.4. (i) *Let $r \geq 1$. Then, for any $m \geq c_r = 4r + 2$, we have that all sets $T^{-i}\mathcal{T}_m$, $i = 0, 1, \dots, r$, are convex. Moreover,*

$$T^{-1}\mathcal{T}_m \subset \mathcal{T}_1, \quad \bigcup_{i=2}^r T^{-i}\mathcal{T}_m \subset \mathcal{T}_2,$$

and, for all $(x, y) \in \mathcal{T}_m$ and $i \in \{1, 2, \dots, r\}$,

$$T^{-i}(x, y) = (x - iy, x - (i - 1)y).$$

(ii) *For any $m \geq c_r$,*

$$T\mathcal{T}_m \subset \mathcal{T}_1, \quad \bigcup_{i=2}^r T^i\mathcal{T}_m \subset \mathcal{T}_2,$$

and, for all $(x, y) \in \mathcal{T}_m$ and $i \in \{2, \dots, r\}$,

$$T^i(x, y) = ((m + 2 - i)y - x, (m + 1 - i)y - x).$$

(iii) Let $j \in \{1, \dots, r\}$. Then

$$\text{length}(\partial T^{j-1} \mathcal{T}_{k_1, \dots, k_r}) \ll_r \frac{1}{k_j}$$

uniformly in $k_1, \dots, k_{j-1}, k_{j+1}, \dots, k_r$ as $k_j \rightarrow \infty$.

Proof. (i) In the beginning we follow closely the proof of Lemma 5 in [3]. The inverse of the transformation T is given by

$$(3.4) \quad T^{-1}(x, y) = \left(\left[\frac{1+y}{x} \right] x - y, x \right), \quad (x, y) \in \mathcal{T}.$$

Since $0 \leq 1 - y < x$, we also have $\left[\frac{1-y}{x} \right] = 0$ and thus, for all $(x, y) \in \mathcal{T}$,

$$(3.5) \quad \kappa(T^{-1}(x, y)) = \left[\frac{1 + \left[\frac{1+y}{x} \right] x - y}{x} \right] = \left[\frac{1+y}{x} \right] + \left[\frac{1-y}{x} \right] = \left[\frac{1+y}{x} \right].$$

Consider next a fixed element $(x, y) \in \mathcal{T}_m$ with $m \geq c_r$. Since $m \geq 5$, we have

$$m \leq \frac{1+x}{y} < m+1 \quad \text{and} \quad x > \frac{m-1}{m+1}.$$

This leads to

$$1 < \frac{1+y}{x} \leq \frac{1 + \frac{1+x}{m}}{x} = \frac{x+m+1}{mx} < \frac{\frac{m-1}{m+1} + m + 1}{m \cdot \frac{m-1}{m+1}} = \frac{m+3}{m-1} \leq 2,$$

showing that $\kappa(T^{-1}(x, y)) = 1$, and - using also (3.4) - that

$$(3.6) \quad T^{-1}(x, y) = (x - y, x) \in \mathcal{T}_1.$$

Next, the inequality $m \geq c_r$ gives

$$(3.7) \quad 1 + \frac{2(2r-1)}{m} \leq 1 + \frac{m-4}{m} \leq 1 + \frac{m-3}{m+1} = \frac{2(m-1)}{m+1}.$$

Thus the inequalities $x > \frac{m-1}{m+1}$ and $y \leq \frac{2}{m}$, fulfilled by $(x, y) \in \mathcal{T}_m$ (see [3, Figure 1]), imply in conjunction with (3.7) that $2x > 1 + (2i-1)y$ for all $(x, y) \in \mathcal{T}_m$ and $i \in \{2, \dots, r\}$, or equivalently that

$$\frac{1+y}{x - (i-1)y} < 2 \quad i \in \{2, \dots, r\}.$$

At the same time, it is clear that $\frac{1+y}{x - (i-1)y} > 1$, so that

$$(3.8) \quad \left[\frac{1+x - (i-2)y}{x - (i-1)y} \right] = 1 + \left[\frac{1+y}{x - (i-1)y} \right] = 2, \quad i \in \{2, \dots, r\}.$$

For $i = 2$, equalities (3.5), (3.7) and (3.8) give

$$\begin{aligned} \kappa(T^{-2}(x, y)) &= \left[\frac{1+x}{x-y} \right] = 2, \\ T^{-2}(x, y) &= (2(x-y) - x, x-y) = (x-2y, x-y), \end{aligned}$$

thus, by (3.5) and by (3.8) with $i = 3$ we have

$$\begin{aligned}\kappa(T^{-3}(x, y)) &= \left\lceil \frac{1+x-y}{x-2y} \right\rceil = 2 \quad \text{and} \\ T^{-3}(x, y) &= (2(x-2y) - x + y, x-2y) = (x-3y, x-2y).\end{aligned}$$

Arguing by induction, it follows at once that, for all $i \in \{2, \dots, r\}$,

$$\begin{aligned}\kappa(T^{-i}(x, y)) &= \left\lceil \frac{1+x-(i-2)y}{x-(i-1)y} \right\rceil = 2 \quad \text{and} \\ T^{-i}(x, y) &= (x-iy, x-(i-1)y).\end{aligned}$$

As a consequence, $T^{-i}\mathcal{T}_m$ is the quadrangle with vertices at $(1 - \frac{2i}{m}, 1 - \frac{2(i-1)}{m})$, $(1 - \frac{2i}{m+1}, 1 - \frac{2(i-1)}{m+1})$, $(1 - \frac{2(i+1)}{m+2}, 1 - \frac{2i}{m+2})$, and $(1 - \frac{2(i+1)}{m+1}, 1 - \frac{2i}{m+1})$. This quadrangle is obviously contained in $\mathcal{T}_2$.

(ii) Let $(x, y) \in \mathcal{T}_m$. Then $T(x, y) = (y, my - x)$, and so

$$\kappa(T(x, y)) = \left\lceil \frac{1+y}{my-x} \right\rceil \geq 1.$$

Since $m \leq \frac{1+x}{y} < m+1$ and $y \leq \frac{2}{m} \leq \frac{1}{3}$, it follows that $(2m-1)y \geq 1 + (2m+2)y - 2 > 1 + 2x$. This leads to $\frac{1+y}{my-x} < 2$, and so we obtain $\kappa(T(x, y)) = 1$. Therefore,

$$T^2(x, y) = (my - x, (m-1)y - x).$$

On the other hand, $y \leq \frac{1+x}{m} < \frac{1+x}{m-i}$; whence

$$(3.9) \quad 2 \leq \left\lceil \frac{1+(m+2-i)y-x}{(m+1-i)y-x} \right\rceil = 1 + \left\lceil \frac{1+y}{(m+1-i)y-x} \right\rceil, \quad i \geq 1.$$

The inequality $m \geq 4r+2$ leads to $m-2r \geq (2r+1)x$, which is equivalent to $(m+1)(1+2x) \leq (2m+1-2r)(1+x)$. Since $1+x < (m+1)y$, we infer that $1+2x < (2m+1-2r)y \leq (2m+1-2i)y$. That is,

$$(3.10) \quad \frac{1+y}{(m+1-i)y-x} < 2, \quad i \in \{1, \dots, r\}.$$

By (3.9) and (3.10), we gather that

$$(3.11) \quad \left\lceil \frac{1+(m+2-i)y-x}{(m+1-i)y-x} \right\rceil = 2, \quad i \in \{2, \dots, r\}.$$

Now we infer inductively that $T^i(x, y) \in \mathcal{T}_2$, and that

$$T^i(x, y) = ((m+2-i)y-x, (m+1-i)y-x), \quad i \in \{2, \dots, r\}.$$

(iii) We use the fact that if Ω_1 and Ω_2 are convex polygons with $\Omega_1 \subseteq \Omega_2$, then $\text{length}(\partial\Omega_1) \leq \text{length}(\partial\Omega_2)$. For $k_j > c_r$ this yields, in conjunction with (i) and (ii),

$$\text{length}(\partial T^{j-1}\mathcal{T}_{k_1, \dots, k_r}) \leq \text{length}(\partial \mathcal{T}_{k_j}) \ll_r \frac{1}{k_j}$$

uniformly in $k_1, \dots, k_{j-1}, k_{j+1}, \dots, k_r$. □

Remark 3.5. Suppose that $(x, y) \in \mathcal{T}_m$ with $m \geq 3$. Then $\frac{1+x}{y} < m+1$ and $y \leq \frac{2}{m}$; hence $\frac{1+y}{my-x} < \frac{1+2/m}{1-y} \leq \frac{1+2/m}{1-2/m} = \frac{m+2}{m-2}$ and thus

$$\bigcup_{m \geq 6} T\mathcal{T}_m \subset \mathcal{T}_1, \quad T(\mathcal{T}_4 \cup \mathcal{T}_5) \subset \mathcal{T}_1 \cup \mathcal{T}_2 \quad \text{and} \quad T\mathcal{T}_3 \subset \mathcal{T}_1 \cup \mathcal{T}_2 \cup \mathcal{T}_3 \cup \mathcal{T}_4.$$

If $(x, y) \in \mathcal{T}_2$, then $y > \frac{1+x}{3} \geq \frac{x}{2} + \frac{1}{6} \geq \frac{1}{3} + \frac{1}{6} = \frac{1}{2}$, and so

$$T\mathcal{T}_2 \subset \mathcal{T}_1 \cup \mathcal{T}_2 \cup \mathcal{T}_3.$$

On the other hand, if $(x, y) \in \mathcal{T}_m$, $m \geq 2$, then it follows by the proof of Lemma 3.4 (i), that $\kappa(T^{-1}(x, y)) < \frac{m+3}{m-1}$. Therefore,

$$\bigcup_{m \geq 5} T\mathcal{T}_m \subset \mathcal{T}_1, \quad T(\mathcal{T}_3 \cup \mathcal{T}_4) \subset \mathcal{T}_1 \cup \mathcal{T}_2 \quad \text{and} \quad T\mathcal{T}_2 \subset \mathcal{T}_1 \cup \mathcal{T}_2 \cup \mathcal{T}_3 \cup \mathcal{T}_4.$$

Owing to the presence of the term R in $C_{R,\Omega}$, we need one more fact, already noticed (in a different form) in [4].

Lemma 3.6. *Let $k \in \mathbb{N}^*$ and let $\mathcal{D}$ be a subset of $\mathcal{T}$. Then the following equalities hold.*

(i) *For k even:*

$$\begin{aligned} N_{\text{odd,even}}(Q(\mathcal{T}_k \cap \mathcal{D})) &= N_{\text{even,odd}}(QT(\mathcal{T}_k \cap \mathcal{D})). \\ N_{\text{even,odd}}(Q(\mathcal{T}_k \cap \mathcal{D})) &= N_{\text{odd,even}}(QT(\mathcal{T}_k \cap \mathcal{D})). \\ N_{\text{odd,odd}}(Q(\mathcal{T}_k \cap \mathcal{D})) &= N_{\text{odd,odd}}(QT(\mathcal{T}_k \cap \mathcal{D})). \end{aligned}$$

(ii) *For k odd:*

$$\begin{aligned} N_{\text{odd,even}}(Q(\mathcal{T}_k \cap \mathcal{D})) &= N_{\text{even,odd}}(QT(\mathcal{T}_k \cap \mathcal{D})). \\ N_{\text{even,odd}}(Q(\mathcal{T}_k \cap \mathcal{D})) &= N_{\text{odd,odd}}(QT(\mathcal{T}_k \cap \mathcal{D})). \\ N_{\text{odd,odd}}(Q(\mathcal{T}_k \cap \mathcal{D})) &= N_{\text{odd,even}}(QT(\mathcal{T}_k \cap \mathcal{D})). \end{aligned}$$

Proof. We denote by T_k the linear transformation defined on $\mathbb{R}^2$ by $T_k(x, y) = (y, ky - x)$. Assume that k is even and let $(a, b) \in Q(\mathcal{T}_k \cap \mathcal{D})$. Then $T(\frac{a}{Q}, \frac{b}{Q}) = (\frac{b}{Q}, \frac{kb}{Q} - \frac{a}{Q})$, so

$$QT(\mathcal{T}_k \cap \mathcal{D}) = \{(b, kb - a); (a, b) \in Q\mathcal{T}_k \cap Q\mathcal{D}\} = T_k(Q(\mathcal{T}_k \cap \mathcal{D})).$$

Moreover, since the matrix that defines T_k is unimodular, the elements of $\mathbb{Z}_{\text{pr}}^2 \cap Q(\mathcal{T}_k \cap \mathcal{D})$ are in 1-1 correspondence with the elements of $\mathbb{Z}_{\text{pr}}^2 \cap T_k(Q(\mathcal{T}_k \cap \mathcal{D})) = \mathbb{Z}_{\text{pr}}^2 \cap Q(T(\mathcal{T}_k \cap \mathcal{D}))$. Besides, we see that a is odd and b is even if and only if b is even and $kb - a$ is odd, implying that

$$\begin{aligned} \#\{(a, b) \in \mathbb{Z}_{\text{pr}}^2 \cap Q(\mathcal{T}_k \cap \mathcal{D}); a \text{ odd}, b \text{ even}\} \\ = \#\{(c, d) \in \mathbb{Z}_{\text{pr}}^2 \cap QT(\mathcal{T}_k \cap \mathcal{D}); c \text{ even}, d \text{ odd}\}. \end{aligned}$$

The other five equalities follow in a similar way. $\square$

Proof of Theorem 1.1. We wish to apply Corollary 3.2 to $\Omega = Q\mathcal{T}_{k_1, \dots, k_r}$. Note first that, since T is area-preserving, we have

$$\text{Area}(\mathcal{T}_{k_1, \dots, k_r}) \leq \text{Area}(T^{-j+1}\mathcal{T}_{k_j}) = \text{Area}(\mathcal{T}_{k_j}) \ll \frac{1}{k_j^3} \quad j \in \{1, \dots, r\}.$$

We claim that for every $j \in \{1, \dots, r\}$, all the numbers $N_{\text{odd,odd}}(Q\mathcal{T}_{k_1, \dots, k_r})$, $N_{\text{odd,even}}(Q\mathcal{T}_{k_1, \dots, k_r})$, and $N_{\text{even,odd}}(Q\mathcal{T}_{k_1, \dots, k_r})$ can be expressed as

$$(3.12) \quad \frac{2Q^2}{\pi^2} \text{Area}(\mathcal{T}_{k_1, \dots, k_r}) + O_r\left(\frac{Q}{k_j} \log Q\right)$$

uniformly in $k_1, \dots, k_{j-1}, k_{j+1}, \dots, k_r$ as $Q \rightarrow \infty$.

If $j \geq 2$, we apply Lemma 3.6 successively $j-1$ times: to k_1 and $\mathcal{D} = T^{-1}\mathcal{T}_{k_2, \dots, k_r}$; to k_2 and $\mathcal{D} = T\mathcal{T}_{k_1} \cap T^{-1}\mathcal{T}_{k_3, \dots, k_r}$; and to k_{j-1} and $\mathcal{D} = T^{j-2}\mathcal{T}_{k_1, \dots, k_{j-2}} \cap T^{-1}\mathcal{T}_{k_j, \dots, k_r}$. This yields

$$N_{\text{odd,odd}}(Q\mathcal{T}_{k_1, \dots, k_r}) = N_{\delta_1, \delta_2}(QT^{j-1}\mathcal{T}_{k_1, \dots, k_r})$$

for some pair $(\delta_1, \delta_2) \in \{(\text{odd}, \text{odd}), (\text{odd}, \text{even}), (\text{even}, \text{odd})\}$ that depends on $k_1, \dots, k_{j-1}$. We may now apply Corollary 3.2 to $\Omega = QT^{j-1}\mathcal{T}_{k_1, \dots, k_r} \subseteq Q\mathcal{T}_{k_j} \subset [0, Q] \times [0, \frac{2Q}{k_j}]$, with

$R \asymp \frac{Q}{k_j}$, $\text{Area}(\Omega) \leq \text{Area}(Q\mathcal{T}_{k_j}) \ll \frac{Q^2}{k_j^3}$, and (according to Lemma 3.4) $\text{length}(\partial\Omega) \ll_r \frac{Q}{k_j}$.

Therefore, we gather that $N_{\text{odd,odd}}(Q\mathcal{T}_{k_1, \dots, k_r})$ is indeed given by (3.12). The same estimates are proved for $N_{\text{odd,even}}(Q\mathcal{T}_{k_1, \dots, k_r})$ and $N_{\text{even,odd}}(Q\mathcal{T}_{k_1, \dots, k_r})$ in a similar fashion.

We may now complete the proof of Theorem 1.1. If $k_j \geq c_r$, then we infer from Lemma 3.4 (i) that $\mathcal{T}_{k_1, \dots, k_r} = \emptyset$ unless $k_1 = \dots = k_{j-2} = k_{j+2} = \dots = k_r = 2$ and $k_{j-1} = k_{j+1} = 1$. On the other hand, we see from [4, Remark 2.3] that $Q\mathcal{T}_{k_1, \dots, k_r} \cap \mathbb{Z}^2 = \emptyset$ unless $\max(k_1, \dots, k_r) \leq 2Q$.

As a result, the only non-zero terms that may appear in the sum from (2.2) arise from paths w having all labels $k_j \leq 2Q$ and at most one $> c_{2h-1}$. Taking now also into account (3.12), the sum $\sum_{w \in \mathfrak{L}_h \cap \mathfrak{S}_\Delta} N_{\text{odd,o}}(v_1)(Q\mathcal{T}_{k_1, \dots, k_{|w|-1}})$ can be expressed as

$$(3.13) \quad \begin{aligned} & \frac{2Q^2}{\pi^2} \sum_{w \in \mathfrak{L}_h \cap \mathfrak{S}_\Delta} \text{Area}(\mathcal{T}_{k_1, \dots, k_{|w|-1}}) + O_h\left(\sum_{k=1}^{2Q} \frac{Q \log Q}{k}\right) \\ &= \frac{2Q^2}{\pi^2} \sum_{w \in \mathfrak{L}_h \cap \mathfrak{S}_\Delta} \text{Area}(\mathcal{T}_{k_1, \dots, k_{|w|-1}}) + O_h(Q \log^2 Q). \end{aligned}$$

The statement in Theorem 1.1 now follows from Proposition 2.1, (3.13), and (1.1). $\square$

4. CONSECUTIVE FAREY FRACTIONS WITH ODD DENOMINATORS IN SHORT INTERVALS

For each interval $I \subseteq [0, 1]$, and each subset $\Omega \subseteq \mathbb{R}^2$, we set

$$\Omega^I = \{(a, b) \in \Omega \cap \mathbb{Z}_{\text{pr}}^2; \bar{b} \in I_a\},$$

where $\bar{b}$ denotes the unique number in $\{1, \dots, a-1\}$ for which $b\bar{b} \equiv 1 \pmod{a}$. If $I = [\alpha, \beta]$, then we also set $I_a = [a(1-\beta), a(1-\alpha)]$.

For any function f defined on Ω , denote

$$S_{f, \text{odd,odd/even}}^I(\Omega) = \sum_{\substack{(a,b) \in \Omega \cap \mathbb{Z}_{\text{pr}}^2 \\ a \text{ odd}, b \text{ odd/even} \\ \bar{b} \in I_a}} f(a, b).$$

The following analog of Proposition 2.1 holds and is similarly proved.

Proposition 4.1. *Let $h \geq 1$, and let $\Delta = (\Delta_1, \dots, \Delta_h) \in (\mathbb{N}^*)^h$. Then, for any interval $I \subseteq [0, 1]$,*

$$N_{Q, \text{odd}}^I(\Delta) = \sum_{w \in \mathfrak{L}_h \cap \mathfrak{S}_\Delta} N_{\text{odd}, o(v_1)}((Q\mathcal{T}_{k_1, \dots, k_{|w|-1}})^I).$$

Proposition 4.2. *Assume that $\Omega \subseteq [0, R_1] \times [0, R_2]$ is a convex region, and that f is a C^1 function on Ω . Then $S_{f, \text{odd}, \text{odd}/\text{even}}^I(\Omega)$ is given by*

$$|I|S'_{f, \text{odd}, \text{odd}/\text{even}}(\Omega) + O_\varepsilon(\|f\|_{L^\infty(\Omega)}(R_2 \log R_1 + m_f R_1^{1/2+\varepsilon}(R_1 + R_2)))$$

for every $\varepsilon > 0$, where m_f is an upper limit for the number of intervals of monotonicity of the functions $y \mapsto f(x, y)$.

Proof. The proof is similar to that of Lemma 8 in [3]. As in [3, (65)], we write

$$(4.1) \quad S_{f, \text{odd}, \text{odd}/\text{even}}^I(\Omega) = S_1 + S_2,$$

where

$$(4.2) \quad \begin{aligned} S_1 &= \sum_{\substack{(a,b) \in \Omega \cap \mathbb{Z}_{\text{pr}}^2 \\ a \text{ odd}, b \text{ odd/even}}} f(a, b) \sum_{x \in I_a} \frac{1}{a} \\ &= \sum_{\substack{(a,b) \in \Omega \cap \mathbb{Z}_{\text{pr}}^2 \\ a \text{ odd}, b \text{ odd/even}}} f(a, b) \frac{1}{a} (|I_a| + O(1)) \\ &= |I|S'_{f, \text{odd}, \text{odd}/\text{even}}(\Omega) + O(\|f\|_{L^\infty(\Omega)} R_2 \log R_1) \end{aligned}$$

and

$$S_2 = \sum_{\substack{(a,b) \in \Omega \cap \mathbb{Z}_{\text{pr}}^2 \\ a \text{ odd}, b \text{ odd/even}}} f(a, b) \sum_{x \in I_a} \frac{1}{a} \sum_{l=1}^{a-1} e\left(\frac{l(\bar{b} - x)}{a}\right).$$

As in [3, (67)], we write

$$(4.3) \quad S_2 = \sum_{\substack{a \in \text{pr}_1(\Omega) \\ a \text{ odd}}} \frac{1}{a} \sum_{l=1}^{a-1} \left(\sum_{x \in I_a} e\left(-\frac{lx}{a}\right) \right) S_{f, \text{odd}/\text{even}, I'_a}(l, a),$$

where $I'_a = \{b; (a, b) \in \Omega\}$ is an interval for every a in the projection $\text{pr}_1(\Omega)$ of Ω on the first coordinate. Here, for any interval J , we denote

$$(4.4) \quad S_{f, \text{odd}/\text{even}, J}(l, a) = \sum_{\substack{b \in J \\ b \text{ odd/even} \\ \gcd(a, b)=1}} f(a, b) e\left(\frac{l\bar{b}}{a}\right)$$

and

$$S_{f, J}(l, a) = \sum_{\substack{b \in J \\ \gcd(a, b)=1}} f(a, b) e\left(\frac{l\bar{b}}{a}\right).$$

By [3, Lemma 9] we have

$$(4.5) \quad |S_{f,J}(l, a)| \ll_{\varepsilon} R_{\Omega, f, J, l, a, \varepsilon},$$

where

$$R_{\Omega, f, J, l, a, \varepsilon} = m_f \|f\|_{L^{\infty}(\Omega)} (|J|a^{-1/2+\varepsilon} + a^{1/2+\varepsilon}) \gcd(l, a)^{1/2}.$$

Writing now

$$S_{f, \text{even}, J}(l, a) = \sum_{\substack{c \in J/2 \\ \gcd(a, c)=1}} f(a, 2c) e\left(\frac{2l\bar{c}}{a}\right) = S_{f_2, J/2}(\bar{2}l, a),$$

where $f_2(x, y) = f(x, 2y)$, and then using (4.5) and $S_{f, \text{odd}, J}(l, a) = S_{f, J}(l, a) - S_{f, \text{even}, J}(l, a)$, we infer that

$$(4.6) \quad \max(|S_{f, \text{even}, J}(l, a)|, |S_{f, \text{odd}, J}(l, a)|) \ll_{\varepsilon} R_{\Omega, f, J, l, a, \varepsilon}.$$

As in [3, (67)-(69)], we infer - from (4.3), (4.4), (4.6), from the fact that the inner sum in (4.3) is a geometric progression which is $\ll \left(\frac{a}{l}, \frac{a}{a-l}\right)$, and from $|I'_a| \leq R_2$ - that

$$(4.7) \quad \begin{aligned} |S_2| &\ll \sum_{a=1}^{R_1} \frac{1}{a} \sum_{l=1}^{a-1} \frac{a}{l} |S_{f, \text{odd/even}, I'_a}(l, a)| \\ &\ll_{\varepsilon} m_f \|f\|_{L^{\infty}(\Omega)} R_1^{1/2+\varepsilon} (R_1 + R_2). \end{aligned}$$

The desired conclusion follows now from (4.1), (4.2) and (4.7). $\square$

Corollary 4.3.

$$N_{\text{odd, odd/even}}((Q\mathcal{T}_{k_1, \dots, k_r})^I) = |I| N_{\text{odd, odd/even}}(Q\mathcal{T}_{k_1, \dots, k_r}) + O_{\varepsilon}(Q^{3/2+\varepsilon}).$$

Theorem 1.3 is now a consequence of Proposition 4.1 and Corollary 4.3.

Acknowledgments. We are grateful to the referee for careful reading of the manuscript and pertinent suggestions that led to the improvement of this paper.

REFERENCES

1. V. Augustin, F.P. Boca, C. Cobeli, A. Zaharescu, *The h-spacing distribution between Farey points*, Math. Proc. Camb. Phil. Soc. **131** (2001), 23–38.
2. F.P. Boca, C. Cobeli, A. Zaharescu, *Distribution of lattice points visible from the origin*, Comm. Math. Phys. **213** (2000), 433–470.
3. F.P. Boca, C. Cobeli and A. Zaharescu, *A conjecture of R.R. Hall on Farey points*, J. Reine Angew. Math. **535** (2001), 207–236.
4. F.P. Boca, R.N. Gologan, A. Zaharescu, *On the index of Farey sequences*, preprint arXiv math.NT/0201044, Quart. J. Math. Oxford Ser. (2) **53** (2002), 377–391.
5. T. Esterman, *On Kloosterman's sums*, Mathematika **8** (1961), 83–86.
6. J. Franel, *Les suites de Farey et le problème de nombres premiers*, Göttinger Nachr., 1924, 198–201.
7. R.R. Hall, *A note on Farey series*, J. London Math. Soc. **2** (1970), 139–148.
8. R.R. Hall, *On consecutive Farey arcs II*, Acta Arith. **66** (1994), 1–9.
9. R.R. Hall, G. Tenenbaum, *On consecutive Farey arcs*, Acta Arith. **44** (1984), 397–405.
10. R.R. Hall, P. Shiu, *The index of a Farey sequence*, Michigan Math. J. **51** (2003), 209–223.
11. A. Haynes, *A note on Farey fractions with odd denominators*, J. Number Theory **98** (2003), 89–104.
12. C. Hooley, *An asymptotic formula in the theory of numbers*, Proc. London Math. Soc. **7** (1957), 396–413.
13. M.N. Huxley, *The distribution of Farey points I*, Acta Arith. **18** (1971), 281–287.

14. M.N. Huxley, A. Zhigljavsky, *On the distribution of Farey fractions and hyperbolic lattice points*, Period. Math. Hungarica **42** (2001), 191–198.
15. E. Landau, *Bemerkungen zu der vorstehenden Abhandlung von Herrn Franel*, Göttinger Nachr. 1924, 202–206.
16. A. Weil, *On some exponential sums*, Proc. Nat. Acad. Sci. U.S.A. 34 (1948), 204–207.

FPB AND AZ: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, 1409 W. GREEN STR., URBANA IL 61801, USA

FPB, CC AND AZ: INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, P.O. BOX 1-764, RO-014700 BUCHAREST, ROMANIA

FBOCA@MATH.UIUC.EDU, CRISTIAN.COBELI@IMAR.RO, ZAHARESCU@MATH.UIUC.EDU